

searches. There was a strong positive association between the number of Tweets showing negative ($r = .711, p = .024$) and positive attitude ($r = .703, p = .026$) and time in years.

Conclusions: Over one-third of Nexplanon-related Tweets reported perceived side effects reportedly associated with Nexplanon which may be of concern AYA patients. While Tweets regarding Nexplanon positively may become more popular, displays of both positive and negative attitudes toward the implant increase over time. Future studies should include social media content analyses of AYA experiences with other FDA-approved devices and large-scale surveys about concerning devices.

Sources of Support: This project was completed without funding support.

RESEARCH POSTER PRESENTATION I: ADVERSE CHILDHOOD EVENTS

110.

THE ASSOCIATION OF SUBJECTIVE SOCIAL STATUS WITH ADVERSE CHILDHOOD EXPERIENCES AND SOCIOECONOMIC STATUS AMONG HISPANIC ADOLESCENTS

Arjun Deven, B.S.¹, Young-Rock Hong¹, Faith A. Newsome¹, Alexandra M. Lee², Darci R. Miller³, Michelle Cardel¹

¹University of Florida; ²The Pennsylvania State University; ³Moffitt Cancer Center.

Purpose: Subjective Social Status (SSS), or one's self-perceived rank on the social ladder, is a strong contributor to adolescent health and psychosocial wellbeing. Among adolescents, low SSS has been associated with increased risk for obesity, depression, and future substance use. SSS is often assessed in the context of one's community or society, and is paired with traditional measures of socioeconomic status (SES) to capture objective status and subjective perceptions regarding one's trajectory and opportunities in life. However, little is known about the determinants of SSS, particularly the influence of childhood adversity. Adverse Childhood Experiences (ACEs) have lasting effects on physical and mental health in youth, and include experiences of abuse, neglect, and household dysfunction. This study examines the association of society/community SSS with ACEs and SES measures of household income, parental education, and food insecurity.

Methods: Hispanic adolescents ($n=133$; age 15-21; 60.2% female) were recruited via flyers and word-of-mouth. To assess SSS, participants completed validated MacArthur scales through self-ranking on a 10-rung ladder representing their individual position relative to their community, and their family's position relative to society. The number of ACEs were assessed using the Modified ACE Questionnaire, expanded to include experiences such as physical bullying and cyber bullying. SES was measured through self-reported questionnaires that assessed household income, parental education, and food insecurity. Univariate analysis was used to explore participant characteristics, and bivariate Pearson's correlations were used to assess associations of SSS with ACEs and SES measures.

Results: The majority (87.2%) of participants experienced at least 1 ACE, and 28.6% of participants experienced 4 or more ACEs. Roughly a fourth (25.6%) of participants experienced food insecurity. We found that society and community SSS, respectively, were each associated with ACEs ($r = -0.219, p = 0.011$; $r = -0.176, p = 0.042$), food insecurity ($r = -0.228, p = 0.008$; $r = -0.278, p = 0.001$), and household income

($r = 0.224, p = 0.009$; $r = 0.204, p = 0.019$). Society nor community SSS were associated with parental education.

Conclusions: To our knowledge, this is the first study to examine the relationship between SSS and ACEs. We found that higher number of ACEs were associated with poor SSS among Hispanic adolescents. While several explanations for the pathway from ACEs to health risks have previously been proposed in adults, there is little consensus in literature and for youth. Here, we suggest SSS may be a mechanism through which ACEs contribute to these outcomes. In this study, we also found that SSS was associated with food insecurity and household income, but not parental education, which is consistent with prior research. Given that SSS is a strong predictor of health and psychosocial outcomes, characterizing the factors that drive the development of SSS among youth may help identify important clinical targets.

Sources of Support: None.

111.

CHART REVIEW OF LATINX YOUTH PATIENTS IN AQUI PARA TI: PARENTAL SEPARATION AND ADOLESCENT HEALTH AND WELL-BEING

Lauren E. McPherson, MD, MAT¹, Maura Shramko, PhD, MPP¹, Jia-Shyuan K. Su, MD, MS¹, Maria Veronica Svetaz, MD, MPH, FFAFP, FSAHM²

¹University of Minnesota; ²University of Minnesota, Hennepin Healthcare.

Purpose: In the US, 1 in 4 Latinx children belongs to a family with mixed immigration status; at least one parent being undocumented – meaning a significant portion of these children may be separated from a caregiver due to deportation. In 2017, with the increase in anti-immigrant policies and rhetoric, Aqui Para Ti/Here for You (APT), a clinic-based youth development program for Latinx youth and their families, found an increase in both anxiety and parental separation in its patient population. The purpose of this study is to describe qualitatively immigration journey and deportation related familial separation experiences via manual review of narrative notes in electronic health records.

Methods: This is a retrospective study using chart review of patient's clinic notes. We included notes from all patients in Aqui Para Ti aged 10-24 between April 1, 2016 – December 31, 2019, who marked that they had been separated from at least 1 parent on the amplified Guidelines for Adolescent Preventive Services (GAPS) questionnaire. We then trained coders to review clinic notes among adolescents who reported separation to identify different forms of separation and record any information provided about the effect of the separation on the patient. Data analysis includes descriptive statistics to examine prevalence of each form of separation among this patient population. In addition, we will examine bivariate and multivariate associations between specific forms of separation and mental and psychosocial health among all APT patients in this window. Research was determined to be exempt by the Hennepin Healthcare Research Institute IRB.

Results: Initial data show that of the 315 APT patients identified during our study time period, 83 experienced parental separation verified after chart review. On initial review, 45% of patients had been separated from both at parents at some time, 54% of patients had only been separated from their father, and 1% of patients had only been separated from their mother. Of those who experienced separation, over 28% of patients experienced separation as a result of deportation